

SUSPECT ADVERSE REACTION REPORT

I. REACTION INFORMATION

1. PATIENT INITIALS (first, last) PRIVACY	1a. COUNTRY COSTA RICA	2. DATE OF BIRTH			2a. AGE 34 Years	3. SEX Female	3a. WEIGHT Unk	4-6 REACTION ONSET			8-12 CHECK ALL APPROPRIATE TO ADVERSE REACTION <input type="checkbox"/> PATIENT DIED <input type="checkbox"/> INVOLVED OR PROLONGED INPATIENT HOSPITALISATION <input type="checkbox"/> INVOLVED PERSISTENT OR SIGNIFICANT DISABILITY OR INCAPACITY <input type="checkbox"/> LIFE THREATENING <input type="checkbox"/> CONGENITAL ANOMALY <input checked="" type="checkbox"/> OTHER
		Day	Month	Year				Day	Month	Year	
7 + 13 DESCRIBE REACTION(S) (including relevant tests/lab data) Event Verbatim [PREFERRED TERM] (Related symptoms if any separated by commas) Other Serious Criteria: Medically Significant suicidal thoughts [Suicidal ideation] depression [Depression] panic attacks [Panic attack] anxiety increased [Anxiety] abrupt weight loss [Weight decreased] ozempic prescribed for weight loss [Off label use] Ozempic dose administered as 0.75 mg [Incorrect dose administered]											
Case Description: Study ID: 199-NovoDia											(Continued on Additional Information Page)

II. SUSPECT DRUG(S) INFORMATION

14. SUSPECT DRUG(S) (include generic name) #1) Semaglutide B 1.34 mg/ml PDS290 (SEMAGLUTIDE 1.34 mg/mL) Solution for injection, 1.34 mg/mL (Continued on Additional Information Page)		20. DID REACTION ABATE AFTER STOPPING DRUG? <input type="checkbox"/> YES <input type="checkbox"/> NO <input checked="" type="checkbox"/> NA
15. DAILY DOSE(S) #1) 0.25 mg, qw	16. ROUTE(S) OF ADMINISTRATION #1) Subcutaneous	
17. INDICATION(S) FOR USE #1) weight loss (Weight control)		21. DID REACTION REAPPEAR AFTER REINTRODUCTION? <input type="checkbox"/> YES <input type="checkbox"/> NO <input checked="" type="checkbox"/> NA
18. THERAPY DATES(from/to) #1) FEB-2024 / MAY-2024	19. THERAPY DURATION #1) Unknown	

III. CONCOMITANT DRUG(S) AND HISTORY

22. CONCOMITANT DRUG(S) AND DATES OF ADMINISTRATION (exclude those used to treat reaction) #1) ENALAPRIL (ENALAPRIL) Tablet ; Ongoing		
23. OTHER RELEVANT HISTORY. (e.g. diagnostics, allergies, pregnancy with last month of period, etc.) From/To Dates Type of History / Notes Description Unknown to Ongoing Current Condition Anxiety (Anxiety) Unknown to Ongoing Current Condition Blood pressure high (Hypertension)		

IV. MANUFACTURER INFORMATION

24a. NAME AND ADDRESS OF MANUFACTURER Novo Nordisk A/S Lise Grimmeshave Vandtaarnsvej 114 Soeborg, DK-2860 DENMARK Phone: +45 44448888		26. REMARKS Medically Confirmed: No
	24b. MFR CONTROL NO. 1417202	25b. NAME AND ADDRESS OF REPORTER NAME AND ADDRESS WITHHELD.
24c. DATE RECEIVED BY MANUFACTURER 22-APR-2025	24d. REPORT SOURCE <input checked="" type="checkbox"/> STUDY <input type="checkbox"/> LITERATURE <input type="checkbox"/> HEALTH PROFESSIONAL <input type="checkbox"/> OTHER:	
DATE OF THIS REPORT 29-APR-2025	25a. REPORT TYPE <input checked="" type="checkbox"/> INITIAL <input type="checkbox"/> FOLLOWUP:	

29-Apr-2025 12:55

ADDITIONAL INFORMATION**7+13. DESCRIBE REACTION(S) continued**

Study description: Trial Title: Patient support programme to support physician and their daily work to maintain an optimal diabetic control of patients through added value services such as treatment starter kit, nutrition support through NovoDia call center, individual workshops, group workshops and free A1c test.

Patient's height: 159 cm.

Patient's weight and body mass index (BMI) were reported.

This serious Solicited Report from COSTA RICA was reported by a Consumer as "suicidal thoughts(Suicidal ideation)" beginning on 2024 , "depression(Depression)" beginning on 2024 , "panic attacks(Panic attack)" beginning on 2024 , "anxiety increased(Anxiety aggravated)" beginning on 2024 , "abrupt weight loss(Weight loss)" beginning on 2024 , "ozempic prescribed for weight loss(Off label use)" beginning on FEB-2024 , "Ozempic dose administered as 0.75 mg(Inappropriate dose of drug administered)" beginning on 2024 and concerned a 34 Years old Female patient who was treated with Ozempic (SEMAGLUTIDE 1.34 mg/mL) from FEB-2024 and ongoing for "weight loss",

Dosage Regimens:

Ozempic: ??-FEB-2024 to ??-MAY-2024, ??-???-2024 to Not Reported (Dosage Regimen Ongoing);

Current Condition included Anxiety, High blood pressure.

Concomitant medications included ENALAPRIL.

Treatment medications included FLUOXETINE.

On an unknown date in FEB-2024 the ozempic was prescribed for weight loss.

On an unknown date in 2024 the patient experienced depression to the point that she had to seek psychological help and take medications because she was having too many panic attacks. It was after about a month of using the medication that she started experiencing those symptoms, and it got worse when the dose was increased, so she decided to stop because she was not mentally well anymore.

The patient also mentions that although she had anxiety issues before, the use of Ozempic was a trigger (referring to the significant increase in anxiety). She stated that she already knew there were those risks, but she didn't think she would experience them. She continued taking it because she thought it might be temporary, but she reached the dose of 0.75 mg, and that's when she realized she was in a bad place because she literally had suicidal thoughts.

The patient reported that the Ozempic did work for her as she lost a lot of weight(Weight) but experienced a rebound effect afterward because of the depression. She started taking the medications, and they made her happy, and She started eating again' (referring to the rebound effect). The patient was still in a recovery state. It was also a very abrupt weight loss, her food intake decreased significantly.

It was reported that everything came back quadrupled, she don't know why, but she feels that injection really worsened her condition. She always told her psychologist 'The time she was thinnest was the time she was saddest in her entire life,' and she states that despite the effects, she continued the medication until she finished it.

The Batch Numbers for Ozempic was requested.

Action taken to Ozempic was reported as No Change.

The outcome for the event "suicidal thoughts(Suicidal ideation)" was Recovering/resolving.

The outcome for the event "depression(Depression)" was Recovering/resolving.

The outcome for the event "panic attacks(Panic attack)" was Recovering/resolving.

The outcome for the event "anxiety increased(Anxiety aggravated)" was Recovering/resolving.

The outcome for the event "abrupt weight loss(Weight loss)" was Recovering/resolving.

On MAY-2024 the outcome for the event "ozempic prescribed for weight loss(Off label use)" was Recovered.

On 2024 the outcome for the event "Ozempic dose administered as 0.75 mg(Inappropriate dose of drug administered)" was Recovered.

Reporter's causality (Ozempic) -

suicidal thoughts(Suicidal ideation) : Possible

depression(Depression) : Possible

panic attacks(Panic attack) : Possible

anxiety increased(Anxiety aggravated) : Possible

abrupt weight loss(Weight loss) : Possible

ADDITIONAL INFORMATION**7+13. DESCRIBE REACTION(S) continued**

ozempic prescribed for weight loss(Off label use) : Unknown
Ozempic dose administered as 0.75 mg(Inappropriate dose of drug administered) : Unknown

Company's causality (Ozempic) -

suicidal thoughts(Suicidal ideation) : Unlikely
depression(Depression) : Unlikely
panic attacks(Panic attack) : Unlikely
anxiety increased(Anxiety aggravated) : Unlikely
abrupt weight loss(Weight loss) : Possible
ozempic prescribed for weight loss(Off label use) : Possible
Ozempic dose administered as 0.75 mg(Inappropriate dose of drug administered) : Possible

Company comment:

Suicidal ideation, depression, panic attack and anxiety are assessed as unlisted; weight decreased, off label use and incorrect dose administered are assessed as listed events according to the Novo Nordisk current CCDS information on Ozempic. The information on relevant medical history (on traumatic or stressful events, substance abuse) are not available. However, a medical history of anxiety is a confounder in this case. Hence the causality is assessed as unlikely for the events Suicidal ideation, depression, panic attack and anxiety; possible for events weight decreased, off label use and incorrect dose administered. This single case report is not considered to change the current knowledge of the safety profile of Ozempic.

14-19. SUSPECT DRUG(S) continued

14. SUSPECT DRUG(S) (include generic name)	15. DAILY DOSE(S); 16. ROUTE(S) OF ADMIN	17. INDICATION(S) FOR USE	18. THERAPY DATES (from/to); 19. THERAPY DURATION
#1) Semaglutide B 1.34 mg/ml PDS290 (SEMAGLUTIDE 1.34 mg/mL) Solution for injection, 1.34 mg/mL; Regimen #2	0.75 mg, qw; Subcutaneous	weight loss (Weight control)	2024 / Ongoing; Unknown