

# SUSPECT ADVERSE REACTION REPORT

## I. REACTION INFORMATION

1. PATIENT INITIALS (first, last) <b>PRIVACY</b>	1a. COUNTRY <b>COSTA RICA</b>	2. DATE OF BIRTH			2a. AGE <b>58</b> Years	3. SEX <b>Female</b>	3a. WEIGHT <b>75.00</b> kg	4-6 REACTION ONSET			8-12 CHECK ALL APPROPRIATE TO ADVERSE REACTION  <input type="checkbox"/> PATIENT DIED  <input type="checkbox"/> INVOLVED OR PROLONGED INPATIENT HOSPITALISATION <input type="checkbox"/> INVOLVED PERSISTENT OR SIGNIFICANT DISABILITY OR INCAPACITY  <input type="checkbox"/> LIFE THREATENING <input type="checkbox"/> CONGENITAL ANOMALY <input checked="" type="checkbox"/> OTHER	
		Day	Month	Year			Day	Month	Year			
										<b>PRIVACY</b>	<b>MAY</b>	<b>2025</b>

7 + 13 DESCRIBE REACTION(S) (including relevant tests/lab data)  
 Event Verbatim [PREFERRED TERM] (Related symptoms if any separated by commas)  
**Other Serious Criteria: Medically Significant**  
**intestinal paralysis [Ileus paralytic]**  
**feels quite confused [Confusional state]**  
**vomits [Vomiting]**  
**feel unwell [Malaise]**  
**headaches [Headache]**  
**did not decreased her appetite initially [Increased appetite]**  
**treatment had no effect on patient initially [Drug ineffective]**  
**Patient not following the physician's instructions [Treatment noncompliance]**  
 (Continued on Additional Information Page)

## II. SUSPECT DRUG(S) INFORMATION

14. SUSPECT DRUG(S) (include generic name) <b>#1 ) Saxenda (liraglutide 6 mg/mL) Solution for injection, 6 mg/mL</b> (Continued on Additional Information Page)		20. DID REACTION ABATE AFTER STOPPING DRUG?  <input type="checkbox"/> YES <input type="checkbox"/> NO <input checked="" type="checkbox"/> NA
15. DAILY DOSE(S) <b>#1 ) UNK</b>	16. ROUTE(S) OF ADMINISTRATION <b>#1 ) Subcutaneous</b>	
17. INDICATION(S) FOR USE <b>#1 ) For weight loss (Weight control)</b>		21. DID REACTION REAPPEAR AFTER REINTRODUCTION?  <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NA
18. THERAPY DATES(from/to) <b>#1 ) MAY-2025 / Unknown</b>	19. THERAPY DURATION <b>#1 ) Unknown</b>	

## III. CONCOMITANT DRUG(S) AND HISTORY

22. CONCOMITANT DRUG(S) AND DATES OF ADMINISTRATION (exclude those used to treat reaction) <b>#1 ) LEVOTHYROXINE (LEVOTHYROXINE) ; MAY-2025 / Ongoing</b>	
23. OTHER RELEVANT HISTORY. (e.g. diagnostics, allergies, pregnancy with last month of period, etc.) From/To Dates      Type of History / Notes      Description <b>Unknown to Ongoing</b> <b>Current Condition</b> <b>Hypothyroidism (Hypothyroidism)</b>	

## IV. MANUFACTURER INFORMATION

24a. NAME AND ADDRESS OF MANUFACTURER <b>Novo Nordisk A/S</b> <b>Lise Grimmeshave</b> <b>Vandtaarnsvej 114</b> <b>Soeborg, DK-2860 DENMARK</b> <b>Phone: +45 44448888</b>		26. REMARKS <b>Medically Confirmed: No</b>
	24b. MFR CONTROL NO. <b>1456582</b>	25b. NAME AND ADDRESS OF REPORTER <b>NAME AND ADDRESS WITHHELD.</b>
24c. DATE RECEIVED BY MANUFACTURER <b>11-JUN-2025</b>	24d. REPORT SOURCE <input checked="" type="checkbox"/> STUDY <input type="checkbox"/> LITERATURE <input type="checkbox"/> HEALTH PROFESSIONAL <input type="checkbox"/> OTHER:	
DATE OF THIS REPORT <b>20-JUN-2025</b>	25a. REPORT TYPE <input checked="" type="checkbox"/> INITIAL <input type="checkbox"/> FOLLOWUP:	

---

**ADDITIONAL INFORMATION****7+13. DESCRIBE REACTION(S) continued**

she decided to increase the dosage, essentially jumping ahead about 2 weeks [Inappropriate schedule of product administration]

Case Description: Study ID: 828652-My Healthy Journey

Study description: Trial title: This is a 40 weeks digital patient support program with focus on exercise, motivation, nutrition & maintaining strategies (only for patients under Liraglutide 3.0 mg).

Patient's height: 156 cm.

Patient's weight: 75 kg.

Patient's BMI: 30.81854040.

This serious Solicited Report from COSTA RICA was reported by a Consumer as "intestinal paralysis(Paralysis intestinal)" beginning on 03-JUN-2025 , "feels quite confused(Confused)" beginning on 03-JUN-2025 , "vomits(Vomiting)" beginning on 03-JUN-2025 , "feel unwell(Feeling unwell)" beginning on 03-JUN-2025 , "headaches(Headache)" beginning on 03-JUN-2025 , "did not decreased her appetite initially(Increased appetite)" beginning on MAY-2025 , "treatment had no effect on patient initially(Lack of drug effect)" beginning on MAY-2025 , "Patient not following the physician's instructions(Treatment noncompliance)" beginning on 2025 , "she decided to increase the dosage, essentially jumping ahead about 2 weeks(Inappropriate schedule of drug administration)" beginning on 2025 and concerned a 58 Years old Female patient who was treated with Saxenda (liraglutide 6 mg/mL) from MAY-2025 to JUN-2025 for "For weight loss",

**Dosage Regimens:**

Saxenda: ??-MAY-2025 to Not Reported, Not Reported to Not Reported, Not Reported to Not Reported, Not Reported to ??-JUN-2025;

Current Condition: Hypothyroidism.

Concomitant medications included - LEVOTHYROXINE.

Treatment medications included - ELECTROLYTE SOLUTIONS [ELECTROLYTES NOS], ALKA-SELTZER(ACETYLSALICYLIC ACID), AVAMIGRAN(Non codable)

On an unspecified date in MAY 2025, The patient indicates that she was using the treatment saxenda ,but initially it had no effect and did not decreased her appetite, so the patient in-creased the dosage a little, which caused very unpleasant side effects. Patient acknowledges that it was her fault for not following the physician's instructions, and now she feels reluctant to continue.

On 03-JUN-2025 ,the patient experiences intestinal paralysis, vomits, and feels quite con-fused. She believes she will not use it anymore, as she also experiences symptoms like head-aches.When she administers the medication, she starts vomiting 8 hours later, noting that this occurred with all dosages, and when she reached the 2.4 mg dosage, she could not tolerate the symptoms. She did this without the doctor's approval and mentions that after reaching the peak at 8 hours, all symptoms resolve.Additionally, she mentions that Saxenda did not decreased her appetite initially, but after increasing the dose, it did decreased her appetite. However, she expe-rienced symptoms such as vomiting and intestinal paralysis, so she reduced the dosage to 1.8 mg. She paused the treatment for approximately 4 days and resumed it the following week but could not tolerate the symptoms and discontinued it. She also indicates that increasing the dos-age was what made her feel unwell; she was supposed to increase it gradually. Since it did not have much effect on her initially, she decided to increase the dosage and skipped the week that the doctor advised her to wait, essentially jumping ahead about 2 weeks.

**Batch Numbers:**

Saxenda: was requested

Action taken to Saxenda was reported as Product discontinued due to AE.

On 10-JUN-2025 the outcome for the event "intestinal paralysis(Paralysis intestinal)" was Recovered.

On 10-JUN-2025 the outcome for the event "feels quite confused(Confused)" was Recovered.

On 10-JUN-2025 the outcome for the event "vomits(Vomiting)" was Recovered.

On 10-JUN-2025 the outcome for the event "feel unwell(Feeling unwell)" was Recovered.

On 10-JUN-2025 the outcome for the event "headaches(Headache)" was Recovered.

On 2025 the outcome for the event "did not decreased her appetite initially(Increased appetite)" was Recovered.

On 2025 the outcome for the event "treatment had no effect on patient initially(Lack of drug effect)" was Recovered.

On JUN-2025 the outcome for the event "Patient not following the physician's instructions(Treatment noncompliance)" was Recovered.

On JUN-2025 the outcome for the event "she decided to increase the dosage, essentially jumping ahead about 2

**ADDITIONAL INFORMATION****7+13. DESCRIBE REACTION(S) continued**

weeks(Inappropriate schedule of drug administration)" was Recovered.

Reporter's causality (Saxenda) -

intestinal paralysis(Paralysis intestinal) : Possible

feels quite confused(Confused) : Possible

vomits(Vomiting) : Possible

feel unwell(Feeling unwell) : Possible

headaches(Headache) : Possible

did not decreased her appetite initially(Increased appetite) : Possible

treatment had no effect on patient initially(Lack of drug effect) : Possible

Patient not following the physician's instructions(Treatment noncompliance) : Unknown

she decided to increase the dosage, essentially jumping ahead about 2 weeks(Inappropriate schedule of drug administration) : Unknown

Company's causality (Saxenda) -

intestinal paralysis(Paralysis intestinal) : Unlikely

feels quite confused(Confused) : Unlikely

vomits(Vomiting) : Possible

feel unwell(Feeling unwell) : Possible

headaches(Headache) : Possible

did not decreased her appetite initially(Increased appetite) : Unlikely

treatment had no effect on patient initially(Lack of drug effect) : Possible

Patient not following the physician's instructions(Treatment noncompliance) : Possible

she decided to increase the dosage, essentially jumping ahead about 2 weeks(Inappropriate schedule of drug administration) : Possible

Company comment:

Ileus paralytic, confusional state, increased appetite are assessed as unlisted events and malaise, drug ineffective, headache, vomiting, treatment noncompliance are assessed as listed events according to NovoNordisk current reference safety information on Saxenda.

Medical history of hypothyroidism is a significant risk factor for the reported of ileus paralytic in the patient. Considering the safety profile of the suspect, details of treatment received and medical history of hypothyroidism, ileus paralytic is assessed as unlikely related to the suspect.

This single case report is not considered to change the current knowledge of the safety profile of Saxenda.

**14-19. SUSPECT DRUG(S) continued**

14. SUSPECT DRUG(S) (include generic name)	15. DAILY DOSE(S); 16. ROUTE(S) OF ADMIN	17. INDICATION(S) FOR USE	18. THERAPY DATES (from/to); 19. THERAPY DURATION
#1 ) Saxenda (liraglutide 6 mg/mL) Solution for injection, 6 mg/mL; Regimen #2	UNK; Subcutaneous	For weight loss (Weight control)	Unknown; Unknown
#1 ) Saxenda (liraglutide 6 mg/mL) Solution for injection, 6 mg/mL; Regimen #3	2.4 mg, qd; Subcutaneous	For weight loss (Weight control)	Unknown; Unknown
#1 ) Saxenda (liraglutide 6 mg/mL) Solution for injection, 6 mg/mL; Regimen #4	1.8 mg, qd; Subcutaneous	For weight loss (Weight control)	Unknown / JUN-2025; Unknown