| | | | | | | | | | | | | | | | CIC | ΟM | IS F | FOI | RM | | | | |
|---|-------------------------------------|------------|---|----------------|-------------------------|--|--|--------------------------|------|------|---|--|-----------|----------|------------------|----|------|-----|----|--|--|--|--|
| | | | | | | | | | | | | | | | | | | | | | | | |
| SUSPEC | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | П | | _ | П | _ | _ | <u> </u> | _ | Т | Т | Т | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | |
| I. REACTION INFORMATION | | | | | | | | | | | | | | | | | | | | | | | |
| 1. PATIENT INITIALS 1a. COUNTRY 2. DATE OF BIRTH 2a. AGE 3. SEX 3a. WEIGHT 4-6 REACTION ONSET 8-12 CHECK ALL | | | | | | | | | | | | | TO | | | | | | | | | | |
| PRIVACY | PANAMA Day Month Year 51 | | | | | | | emale Unk Day Month Year | | | | | | | ADVERSE REACTION | | | | | | | | |
| 7+13 DESCRIBE REACTION(S) (including relevant tests/lab data) Event Verbatim [LOWER LEVEL TERM] (Related symptoms if any separated by commas) redness (erythema) in the area of the injection [Injection site redness] it was applied intradermally [Incorrect route of product administration] | | | | | | | | | | | | PATIENT DIED INVOLVED OR PROLONGED INPATIENT HOSPITALISATION | | | | | | | | | | | |
| Case Description: | This is a sponta | neous repo | ort received fr | om a Nur | se, Progra | am ID: 164 | 1974. | | | | | _ ا | | | | | | | | | | | |
| A 51-year-old female patient received etanercept (ENBREL), (Lot number: LL4200, Expiration Date: May2027 at 50 mg weekly. | | | | | | | | | | | | 7) INVOLVED PERSISTENT OR SIGNIFICANT DISABILITY OR INCAPACITY | | | | | | | | | | | |
| (Continued on Additional Information Page | | | | | | | | | | age) | LIFE THREATENING | | | | | | | | | | | | |
| | | | I. SUSPEC | T DRU | IG(S) IN | FORMA | TIO | N | | | | | | | | | | | | | | | |
| 14. SUSPECT DRUG(S) (include generic name) #1) Enbrel (ETANERCEPT) Solution for injection in pre-filled syringe {Lot # LL4200; Exp.Dt. MAY-2027} #2) Enbrel (ETANERCEPT (DEVICE CONSTITUENT)) Solution for injection in pre-filled syringe | | | | | | | | | | | 20. DID REACTION ABATE AFTER STOPPING DRUG? | | | | | | | | | | | | |
| 15. DAILY DOSE(S) #1) 50 mg, weekly #2) | | | | | | . ROUTE(S) OF ADMINISTRATION) Unknown 2) Unknown | | | | | | | YES NO NA | | | | | | | | | | |
| 17. INDICATION(S) FOR USE #1) Unknown #2) Unknown | | | | | | | | | | | 21. DID REACTION REAPPEAR AFTER REINTRODUCTION? | | | | | | | | | | | | |
| #1) Unknown #1 | | | | | | THERAPY DURATION) Unknown) Unknown | | | | | | | YES NO NA | | | | | | | | | | |
| #2) Gillatouri | | | CONCOMI | | | | пот | OD: | | | | | | | | | | | | | | | |
| 22. CONCOMITANT DRU | G(S) AND DATES OF A | | N (exclude those us | | , | AND | 11011 | <u>UK</u> | I | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | |
| 23. OTHER RELEVANT H From/To Dates Unknown | IISTORY. (e.g. diagnosti | | egnancy with last m of History / Notes | onth of period | d, etc.) Description | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | |
| | | | IV. MANUF | FACTU | | | ΓΙΟΝ | 1 | | | | | | | | | | | | | | | |
| 24a. NAME AND ADDRES Pfizer S.A. Laura Arce Mora Avenida Escazú, To San jose, COSTA | 26. REM | IARKS | | | | | | | | | | | | | | | | | | | | | |
| | 24b. MFR CONTROL NO. PV202500088178 | | | | | | 25b. NAME AND ADDRESS OF REPORTER NAME AND ADDRESS WITHHELD. | | | | | | | | | | | | | | | | |
| 24c. DATE RECEIVED | 24d. REPC | ORT SOURCE | | | NAME | AND ADD | RES | s WI | THHE | ELD. | | | | | | | | | | | | | |
| 19-JUL-2025 | R STUD | | | | | | | | | | | | | | | | | | | | | | |
| DATE OF THIS REPORT 24-JUL-2025 | 25a. REPO | | FOLLOWUP: | | | | | | | | | | | | | | | | | | | | |

ADDITIONAL INFORMATION

7+13. DESCRIBE REACTION(S) continued

The patient's relevant medical history and concomitant medications were not reported.

The following information was reported: INJECTION SITE ERYTHEMA (non-serious), outcome "recovered", described as "redness (erythema) in the area of the injection"; INCORRECT ROUTE OF PRODUCT ADMINISTRATION (non-serious), outcome "unknown", described as "it was applied intradermally". The action taken for etanercept was unknown.

Additional Information: The nurse indicated that the patient applied her weekly dose, but did not do it by the subcutaneously, it was applied intradermally, which caused a reaction: redness (erythema) in the area of the injection, however, she clarified that the discomfort was temporary.