

SUSPECT ADVERSE REACTION REPORT	
SI-Tolmar-TLM-2025-01726	

I. REACTION INFORMATION

1. PATIENT INITIALS (first, last)	1a. COUNTRY	2. DATE OF BIRTH			2a. AGE Years	3. SEX	4-6 REACTION ONSET			8-12 CHECK ALL APPROPRIATE TO ADVERSE REACTION <input type="checkbox"/> PATIENT DIED <input type="checkbox"/> LIFE THREATENING <input type="checkbox"/> INVOLVED OR PROLONGED INPATIENT HOSPITALIZATION <input type="checkbox"/> RESULTS IN PERSISTENCE OR SIGNIFICANT DISABILITY/INCAPACITY <input type="checkbox"/> CONGENITAL ANOMALY <input type="checkbox"/> OTHER MEDICALLY IMPORTANT CONDITION
CAMN	EL	Day	Month	Year	79	Male	Day	Month	Year	
	Cont..	20	Nov	1945						
7+13 DESCRIBE REACTION(S) (including relevant tests/lab data) 1) Pain in the area of application (Application site pain (10003051), Application site pain (10003051)) Recovered/Resolved										

II. SUSPECT DRUG(S) INFORMATION

14. SUSPECT DRUG(S) (include generic name) 1) Eligard® (Leuprolide acetate, Leuprolide acetate) (Suspect) (22.5 Milligram, Injection) (Unknown)	Cont..	20. DID EVENT ABATE AFTER STOPPING DRUG? <input type="checkbox"/> YES <input type="checkbox"/> NO <input checked="" type="checkbox"/> NA 21. DID EVENT REAPPEAR AFTER REINTRODUCTION <input type="checkbox"/> YES <input type="checkbox"/> NO <input checked="" type="checkbox"/> NA (NA : Not Applicable)	
15. DAILY DOSE(S) 1) (22.5 milligram(s), 1 in 3 Month)	Cont..		
16. ROUTE(S) OF ADMINISTRATION 1) Subcutaneous	Cont..		
17. INDICATION(S) FOR USE 1) Prostate cancer [10060862 - Prostate cancer]			
18. THERAPY DATE(S) (from/to) 1) (18/Mar/2025 - Ongoing)		19. THERAPY DURATION	

III. CONCOMITANT DRUG(S) AND HISTORY

22. CONCOMITANT DRUG(S) AND DATES OF ADMINISTRATION (exclude those used to treat reaction) 1) BENICAR(OLMESARTAN MEDOXOMIL)	Cont..
23. OTHER RELEVANT HISTORY (e.g. diagnostics, allergies, pregnancy with last month of period, etc.) 1) PROSTATE CANCER (10060862, Prostate cancer) (Continuing: Yes)	Cont..

IV. MANUFACTURER INFORMATION

24a. NAME AND ADDRESS OF MANUFACTURER Name : Tolmar, Inc 701 Centre Avenue Fort Collins, CO, 80526, UNITED STATES OF AMERICA debbie.maierhofer@tolmar.comand+1-4129158447		Study Information Study Name: NA EudraCT Number: Protocol No.: NA Center No.: Subject Id :
24. REPORT NULLIFIED <input type="checkbox"/> YES <input type="checkbox"/> NO	24b. MFR CONTROL NO. SI-Tolmar-TLM-2025-01726	
24c. DATE RECEIVED BY MANUFACTURER 15/May/2025	24d. REPORT SOURCE <input checked="" type="checkbox"/> STUDY <input type="checkbox"/> LITERATURE <input type="checkbox"/> HEALTH PROFESSIONAL	
DATE OF THIS REPORT 22/May/2025	25a. REPORT TYPE <input checked="" type="checkbox"/> INITIAL <input type="checkbox"/> FOLLOWUP	

= Continuation attached sheet(s)..

Continuation Sheet for CIOMS report

1a. COUNTRY

EL SALVADOR

7+13 DESCRIBE REACTION(S) (including relevant tests/lab data) (Continuation...)

Event Description :

This study report from El salvador was received by Adium via 'ASOFARMA A TU LADO' Patient Support Program (reference number: SV-ADIUM-SV-0005-20250508) on 08-May-2025 from a consumer (patient's wife) (non-healthcare professional) regarding an elderly 79-year-old male patient who experienced a non-serious event of 'pain in the area of application' (Application site pain) during Eligard (Leuprolide acetate) 45 mg therapy for prostate cancer. The needle component of the constituent device was not identified in the report. The report was sent to Tolmar on 13-May-2025.

The patient's medical history was unknown. Current condition included prostate cancer and hypertension.

Concomitant medication included Benicar (olmesartan medoxomil).

On an unknown date in 2021, the patient began receiving Eligard 45 mg, every 6 months via subcutaneous route for prostate cancer (Lot numbers and Expiration dates were not provided).

On an unknown date, about 2 to 3 months ago the patient received Eligard 45 mg, every 6 months via subcutaneous route for prostate cancer and experienced a lot of pain around the stomach where the Eligard was applied. The patient presented pain only with one dose, while previous doses did not cause pain, and had probably received 8 doses of Eligard.

It was reported that the patient suffered from high blood pressure long before Eligard. According to CRM data, he started Eligard on 02-Apr-2024, but his wife mentioned the treatment began in 2021.

On an unknown date in Jun-2025 (proposed commencement date), patient next application was scheduled.

Corrective treatment was not reported.

Action taken with Eligard in response to the event was dose not changed. De-challenge and re-challenge were not applicable.

On an unknown date, the outcome of application site pain was resolved.

The reporter did not assess the seriousness of application site pain.

The reporter assessed the causality of application site pain in relationship to Eligard and Eligard unspecified device as related.

No further queries were raised.

Note: Since the primary reporter is patient wife and the start date of Eligard was captured as reported by her (Which will match with patient being taken 8 doses of Eligard).

On 15-May-2025, follow up information was received via by Adium via Patient Support Program "ASOFARMA A TU LADO (reference number: (SV-ADIUM-SV-0005-20250508 (1))) from a consumer (non-healthcare professional). On 16-May-2025 additional information was received. Both were processed together. New information included: New Eligard 22.5mg dose details were added. Action taken with Eligard 45 mg was updated from 'dose not changed' to 'unknown'.

On 18-Mar-2025, the patient began receiving Eligard 22.5 mg every 3 months via subcutaneous route for prostate cancer (Lot numbers and Expiration dates were not provided). It was reported that the only dose that caused pain to patient is the one that was applied on this day.

It was unknown if the patient will continue using Eligard 45mg, as he is being followed up by the insurance company, in which he was given with Eligard 22.5 for 3 months.

On an unknown date, in Jun-2025, the patient will be receiving the same dose Eligard 22.5 mg every 3 months.

Action taken with Eligard 22.5 mg, in response to event was dose not changed.

Action taken with Eligard 45 mg in response to event was unknown.

The reporter assessed the causality of application site pain in relationship to Eligard 22.5 mg and 45 mg and Eligard unspecified device as related.

Listedness:

Application site pain>Eligard>listed as per CCDS>07-Nov-2024

Application site pain>Eligard>listed as per USPI>Feb-2025

Application site pain>Eligard unspecified device>listed as per USPI>Feb-2025

Continuation Sheet for CIOMS report

Application site pain>Eligard>listed as per Canadian monograph>02-Apr-2025

Company Remarks (Sender's Comments) :

Evaluator comment (Tolmar): This is regarding an elderly 79-year-old male patient who experienced application site pain (Pain in the area of application), during Eligard (leuprolide acetate) 22.5 mg therapy for prostate cancer. Tolmar assessed the event as non-serious since it does not meet the ICH seriousness criteria and is not an IME event. The reported event application site pain was considered as related to Eligard (drug and device) considering the known pharmacological profile of the drug. The event application site pain was assessed as not related to Eligard 45 mg (drug and device) as the event occurred with 22.5 mg Eligard therapy.

14.SUSPECT DRUG(S) (Continuation...)

Product-Reaction Level

1) Drug : Eligard® (Leuprolide acetate)
 Active Substance : 1) Leuprolide acetate
 Drug Characterization : Suspect
 Form Strength : 1) 22.5 Milligram
 Form of Admin : 1) Injection
 Lot Number : 1) Unknown
 Daily Dose : (22.5 milligram(s), 1 in 3 Month)
 Route of Admin : 1) Subcutaneous
 Indications : 1) Prostate cancer [10060862 - Prostate cancer]
 Therapy Dates : 1) From : 18/Mar/2025 To :Continuing
 Action(s) Taken With Drug : Dose not changed

Causality

1) Pain in the area of application (Application site pain - 10003051, Application site pain - 10003051)
 Causality as per reporter : Related
 Causality as per Mfr : Related
 DeChallenge : Not applicable
 ReChallenge : Not Applicable

Labeling :

1) Pain in the area of application
 CORE Labeled

2) Drug : Eligard® (Leuprolide acetate)
 Active Substance : 1) Leuprolide acetate
 Drug Characterization : Suspect
 Form Strength : 1) 45 Milligram
 Form of Admin : 1) Injection
 Lot Number : 1) Unknown
 Daily Dose : (45 milligram(s), 1 in 6 Month)
 Route of Admin : 1) Subcutaneous
 Indications : 1) Prostate cancer [10060862 - Prostate cancer]
 Therapy Dates : 1) From : //2021 To :Unknown
 Action(s) Taken With Drug : Unknown

Causality

1) Pain in the area of application (Application site pain - 10003051, Application site pain - 10003051)
 Causality as per reporter : Related
 Causality as per Mfr : Not Related
 DeChallenge : Not applicable
 ReChallenge : Not Applicable

Labeling :

1) Pain in the area of application
 CORE Labeled

3) Drug : Eligard® Unspecified Device (Leuprolide acetate)
 Active Substance : 1) Leuprolide acetate
 Drug Characterization : Suspect
 Form of Admin : 1) Injection
 Lot Number : 1) Unknown
 Indications : 1) Prostate cancer [10060862 - Prostate cancer]

Continuation Sheet for CIOMS report

Action(s) Taken With Drug : Not applicable

Causality

1) Pain in the area of application (Application site pain - 10003051, Application site pain - 10003051)

Causality as per reporter : Related

Causality as per Mfr : Related

DeChallenge : Not applicable

ReChallenge : Not Applicable

Labeling :

1) Pain in the area of application

CORE

22.CONCOMITANT DRUG(S) (Continuation...)

1). Drug : BENICAR

Active Substance : 1) OLMESARTAN MEDOXOMIL

Form Strength :

Daily Dose : 1) (1 in 1 Day)

Indications : 1) High blood pressure [10005747 - Blood pressure high]

23. OTHER RELEVANT HISTORY (Continuation...)

2) HIGH BLOOD PRESSURE (10005747 , Blood pressure high) (Continuing : YES)